arXiv:1404.2679v1 [math.MG] 10 Apr 2014

ESTIMATE OF OBSERVABLE DIAMETER OF l_p-PRODUCT SPACES

RYUNOSUKE OZAWA AND TAKASHI SHIOYA

ABSTRACT. We estimate the observable diameter of the l_p-product space X^n of an mm-space X by using the limit formula in our previous paper [9]. The idea of our proof is based on Gromov's book [3, §3.$\frac{1}{2}$]. As a corollary we obtain the phase transition property of $\{X^n\}_{n=1}^{\infty}$ under a discreteness condition.

1. INTRODUCTION

In the study of concentration of measure phenomenon discovered by Lévy and Milman, it is an important problem to estimate the observable diameter (or equivalently the concentration function) of a given mm-space, where an *mm-space* is defined to be a space X equipped with a complete separable metric d_X and a Borel probability measure μ_X on X. The purpose of this paper is to estimate the observable diameter of the *l_p-product space*

$$X_p^n := (X^n, d_{X_p^n}, \mu_X^{\otimes n}),$$

of an mm-space X, where $d_{X_p^n}$ is the *l_p-metric*, $1 \leq p < +\infty$, defined by

$$d_{X_p^n}(x, y) := \left(\sum_{i=1}^{n} d_X(x_i, y_i)^p \right)^{\frac{1}{p}}$$

for $x = (x_1, x_2, \ldots, x_n), y = (y_1, y_2, \ldots, y_n) \in X^n$, and $\mu_X^{\otimes n}$ is the product measure. We refer to [1, 5, 8, 11] for some significant studies of concentration of measure for l_1-product spaces.

One of our main theorems is stated as follows.

Theorem 1.1. *Let X be an mm-space with finite diameter. Then, for any $0 < \kappa < 1$ and $1 \leq p < +\infty$, we have*

$$\mathrm{ObsDiam}(X_p^n; -\kappa) \leq C_{\kappa,p} \, \mathrm{diam}(X) \, n^{\frac{1}{2p}},$$

Date: April 11, 2014.

2010 *Mathematics Subject Classification.* Primary 53C23.

Key words and phrases. metric measure space, product space, concentration of measure, observable diameter, concentration function, phase transition.

The authors are partially supported by a Grant-in-Aid for Scientific Research from the Japan Society for the Promotion of Science. The first author is supported by Research Fellowships of the Japan Society for the Promotion of Science for Young Scientists.

where

$$C_{\kappa,p} := \begin{cases} 4\sqrt{2\log\frac{2}{\kappa}} & \text{if } p = 1, \\ 4 + 4\sqrt{2\log\frac{2}{\kappa}} & \text{if } 0 < p < +\infty. \end{cases}$$

In the case where $p = 1$, Theorem 1.1 is already known (see [5, §1.6]). For $p > 1$, a rough sketch of the proof was given in [3, §3.$\frac{1}{2}$.62]. In this paper, we give a complete proof of the theorem by using the limit formula for observable diameter (see Theorem 2.11) established in our previous paper [9].

For lower estimate, we have the following theorems.

Theorem 1.2. *Let X be a nontrivial mm-space of finite diameter such that $d_X(x, y) \geq \delta$ for any different $x, y \in X$, where δ is a positive constant. Then, for any $0 < \kappa < 1$ and $1 < p < +\infty$, we have*

$$\liminf_{n\to\infty} \frac{\mathrm{ObsDiam}(X_p^n; -\kappa)}{n^{\frac{1}{2p}}} \geq \left(\frac{2\,\Phi^{-1}(\kappa)\sqrt{V_p(X)}}{p}\right)^{\frac{1}{p}} \left(\frac{\delta}{\mathrm{diam}(X)}\right)^{1-\frac{1}{p}},$$

where

$$\Phi(r) := \int_r^\infty \frac{1}{\sqrt{2\pi}} e^{-\frac{t^2}{2}}\, dt, \qquad V_p(X) := \sup_{x\in X} V(d_X(x,\cdot)^p),$$

and $V(f)$ is the variance of a function $f : X \to \mathbb{R}$, i.e., $V(f) := \int_X \left(f - \int_X f d\mu_X\right)^2 d\mu_X$.

If $p = 1$, then the δ-discreteness assumption is not necessary.

Theorem 1.3. *Let X be a nontrivial mm-space of finite diameter. Then, for any $0 < \kappa < 1$, we have*

$$\liminf_{n\to\infty} \frac{\mathrm{ObsDiam}(X_1^n; -\kappa)}{\sqrt{n}} \geq 2\,\Phi^{-1}(\kappa)\sqrt{V_1(X)}.$$

Gromov [3, §3.$\frac{1}{2}$.42] proved Theorem 1.2 for $X = \{0, 1\}$ and $p = 2$.

For Theorem 1.2, the δ-discreteness assumption for X is necessary. In fact, we see from a result of [2] that $\mathrm{ObsDiam}(X_2^n; -\kappa)$ is bounded from above if X is a compact and connected Riemannian manifold (see Proposition 3.4).

In the case where $p = +\infty$, we have the trivial uniform estimates

$$0 < \mathrm{ObsDiam}(X; -\kappa) \leq \mathrm{ObsDiam}(X_\infty^n; -\kappa) \leq \mathrm{diam}(X)$$

for X nontrivial, where $d_{X_\infty^n}(x, y) := \max_{i=1}^n d_X(x_i, y_i)$.

Combining Theorems 1.1, 1.2, and 1.3 tells us that the order of the observable diameter $\mathrm{ObsDiam}(X_p^n; -\kappa)$ as $n \to \infty$ is $n^{\frac{1}{2p}}$, which is independent of κ. This together with Theorem [9, Theorem 1.2] implies the following corollary.

Corollary 1.4. *Let X be a nontrivial mm-space of finite diameter and let $1 \leq p \leq +\infty$. We assume that $\inf_{x,y\in X,\ x\neq y} d_X(x,y) > 0$ if $1 < p < +\infty$. Then, $\{X_p^n\}_{n=1}^{\infty}$ has the phase transition property with critical scale order $\sim n^{-\frac{1}{2p}}$, i.e.,*

(1) *if $n^{\frac{1}{2p}} t_n \to 0$ as $n \to \infty$, then $\{t_n X_p^n\}$ is a Lévy family;*

(2) *if $n^{\frac{1}{2p}} t_n \to +\infty$ as $n \to \infty$, then $\{t_n X_p^n\}$ ∞-dissipates,*

where $t_n X_p^n$ is the scale change of X_p^n of factor t_n.

A Lévy family is a sequence of mm-spaces with observable diameter tending to zero. An ∞-dissipating sequence is an opposite notion to a Lévy family and means that the mm-spaces disperse into many small pieces far apart each other (see [3, 9, 10] for the details). The corollary claims the asymptotic uniformity of the l_p-product spaces X_p^n in a sense.

2. Preliminaries

In this section, we state some definitions and facts needed in this paper. We refer to [3, 10] for more details.

Definition 2.1 (Lipschitz order)**.** Let X and Y be two mm-spaces. We say that X (*Lipschitz*) *dominates* Y and write $Y \prec X$ if there exists a 1-Lipschitz map $f : X \to Y$ with $f_*\mu_X = \mu_Y$, where $f_*\mu_X$ is the push-forward measure of μ_X by f. We call the relation $\prec$ the *Lipschitz order*.

Definition 2.2 (Partial and observable diameter)**.** Let X be an mm-space. For a real number α, we define the *partial diameter* $\operatorname{diam}(X;\alpha) = \operatorname{diam}(\mu_X;\alpha)$ *of* X to be the infimum of $\operatorname{diam}(A)$, where $A \subset X$ runs over all Borel subsets with $\mu_X(A) \geq \alpha$ and $\operatorname{diam}(A)$ denotes the diameter of A. For a real number $\kappa > 0$, we define the *observable diameter of* X to be

$$\operatorname{ObsDiam}(X;-\kappa) := \sup\{\ \operatorname{diam}(f_*\mu_X;1-\kappa) \mid f : X \to \mathbb{R} \text{ is 1-Lipschitz continuous }\}.$$

Clearly, $\operatorname{ObsDiam}(X;-\kappa)$ is monotone nonincreasing in $\kappa > 0$. Note that $\operatorname{ObsDiam}(X;-\kappa) = \operatorname{diam}(X;1-\kappa) = 0$ for $\kappa \geq 1$.

For an mm-space X and a real number $t > 0$, we define tX to be the mm-space X with the scaled metric $d_{tX} := t d_X$.

Proposition 2.3. *Let X be an mm-space. Then we have*

$$\operatorname{ObsDiam}(tX;-\kappa) = t \operatorname{ObsDiam}(X;-\kappa)$$

for any $t, \kappa > 0$.

Proposition 2.4. *If $X \prec Y$, then*

$$\operatorname{ObsDiam}(X;-\kappa) \leq \operatorname{ObsDiam}(Y;-\kappa)$$

for any $\kappa > 0$.

Definition 2.5. The *concentration function* $\alpha_X(r)$, $r > 0$, of an mm-space X is defined by

$$\alpha_X(r) := \sup\{\, 1 - \mu_X(A_r) \mid A \subset X : \text{Borel},\ \mu_X(A) \geq 1/2 \,\},$$

where A_r denotes the open r-neighborhood of A.

Proposition 2.6 (see [5, Proposition 1.12], [10, Remark 2.28])**.** *We have*

$$\text{(1)} \qquad \mathrm{ObsDiam}(X; -\kappa) \leq 2 \inf\{\, r > 0 \mid \alpha_X(r) \leq \kappa/2 \,\},$$

$$\text{(2)} \qquad \alpha_X(r) \leq \sup\{\, \kappa > 0 \mid \mathrm{ObsDiam}(X; -\kappa) \geq r \,\}.$$

Definition 2.7 (Separation distance)**.** Let X be an mm-space. For any real numbers $\kappa_0, \kappa_1, \cdots, \kappa_N > 0$ with $N \geq 1$, we define the *separation distance*

$$\mathrm{Sep}(X; \kappa_0, \kappa_1, \cdots, \kappa_N)$$

of X as the supremum of $\min_{i \neq j} d_X(A_i, A_j)$ over all sequences of $N+1$ Borel subsets $A_0, A_2, \cdots, A_N \subset X$ satisfying that $\mu_X(A_i) \geq \kappa_i$ for all $i = 0, 1, \cdots, N$, where $d_X(A_i, A_j) := \inf_{x \in A_i, y \in A_j} d_X(x, y)$. If there exists no sequence $A_0, \ldots, A_N \subset X$ with $\mu_X(A_i) \geq \kappa_i$, $i = 0, 1, \cdots, N$, then we define

$$\mathrm{Sep}(X; \kappa_0, \kappa_1, \cdots, \kappa_N) := 0.$$

Proposition 2.8. *For any mm-space X and any real numbers κ and κ' with $\kappa > \kappa' > 0$, we have*

$$\mathrm{ObsDiam}(X; -2\kappa) \leq \mathrm{Sep}(X; \kappa, \kappa) \leq \mathrm{ObsDiam}(X; -\kappa').$$

Definition 2.9 (Parameter)**.** Let $I := [\,0, 1\,)$ and let X be an mm-space. A map $\varphi : I \to X$ is called a *parameter of X* if φ is a Borel measurable map such that

$$\varphi_* \mathcal{L}^1 = \mu_X,$$

where $\mathcal{L}^1$ denotes the one-dimensional Lebesgue measure on I.

Any mm-space has a parameter.

Definition 2.10 (Box distance)**.** We define the *box distance* $\square(X, Y)$ *between two mm-spaces X and Y* to be the infimum of $\varepsilon \geq 0$ satisfying that there exist parameters $\varphi : I \to X$, $\psi : I \to Y$, and a Borel subset $I_0 \subset I$ such that

$$\text{(1)} \qquad |\, \varphi^* d_X(s, t) - \psi^* d_Y(s, t) \,| \leq \varepsilon \quad \text{for any } s, t \in I_0;$$

$$\text{(2)} \qquad \mathcal{L}^1(I_0) \geq 1 - \varepsilon,$$

where $\varphi^* d_X(s, t) := d_X(\varphi(s), \varphi(t))$ for $s, t \in I$.

The box distance function $\square$ is a complete separable metric on the set of isomorphism classes of mm-spaces.

The following is a corollary to [9, Theorem 1.1].

Theorem 2.11 (Limit formula for observable diameter; [9])**.** *Let X and X_n, $n = 1, 2, \dots$, be mm-spaces. If X_n $\square$-converges to X as $n \to \infty$, then*

$$\begin{aligned}\operatorname{ObsDiam}(X; -\kappa) &= \lim_{\varepsilon \to 0+} \liminf_{n\to\infty} \operatorname{ObsDiam}(X_n; -(\kappa+\varepsilon)) \\ &= \lim_{\varepsilon \to 0+} \limsup_{n\to\infty} \operatorname{ObsDiam}(X_n; -(\kappa+\varepsilon))\end{aligned}$$

for any $\kappa > 0$.

3. Proof

We need some lemmas for the proof of Theorem 1.1.

Definition 3.1. Let k be a natural number. A *k-regular* mm-space is defined to be a k-point mm-space X satisfying that $d_X(x, y) = 1$ for any different two points $x, y \in X$ and $\mu_X = \frac{1}{k}\sum_{x\in X} \delta_x$, where δ_x denotes the Dirac measure at x.

Lemma 3.2. *Let X be a k-regular mm-space. Then, for any $0 < \kappa < 1$ and $1 \le p < +\infty$, we have*

$$\operatorname{ObsDiam}(X_p^n; -\kappa) \le n^{\frac{1}{2p}-\frac{1}{2}} \operatorname{ObsDiam}(X_1^n; -\kappa) + 4n^{\frac{1}{2p}}.$$

Proof. Let $f : X_p^n \to \mathbb{R}$ be an arbitrary 1-Lipschitz function. For $\delta := n^{\frac{1}{2p}}$, we take a maximal δ-separated net $Y \subset X_p^n$, i.e., Y is a maximal set of X such that any different two points in Y has l_p-distance at least δ. We prove that f is δ^{1-p}-Lipschitz with respect to $d_{X_1^n}$. In fact, since X is k-regular, for any different two points $x, x' \in Y$ we have

$$|f(x) - f(x')| \le d_{X_p^n}(x, x') = d_{X_1^n}(x, x')^{\frac{1}{p}}.$$

So, it suffices to prove that $d_{X_1^n}(x, x')^{\frac{1}{p}} \le \delta^{1-p} d_{X_1^n}(x, x')$, which follows from

$$\delta^{1-p} d_{X_1^n}(x, x')^{1-\frac{1}{p}} = \delta^{1-p} d_{X_p^n}(x, x')^{p-1} \ge \delta^{1-p}\delta^{p-1} = 1,$$

since $d_{X_p^n}(x, x') \ge \delta$.

There is a δ^{1-p}-Lipschitz extension $f' : X_1^n \to \mathbb{R}$ of $f|_Y$. We have

$$\begin{aligned}\delta^{p-1} \operatorname{diam}(f'_* \mu_{X_1^n}; 1-\kappa) &= \operatorname{diam}((\delta^{p-1} f')_* \mu_{X_1^n}; 1-\kappa) \\ &\le \operatorname{ObsDiam}(X_1^n; -\kappa)\end{aligned}$$

and hence

$$\operatorname{diam}(f'_* \mu_{X_1^n}; 1-\kappa) \le n^{\frac{1}{2p}-\frac{1}{2}} \operatorname{ObsDiam}(X_1^n; -\kappa).$$

Let us estimate the distance between f and f'. For any point $x \in X_1^n$ there is a point $x' \in Y$ such that $d_{X_p^n}(x, x') \le \delta$. Then, since

$d_{X_1^n}(x,x') = d_{X_p^n}(x,x')^p \le \delta^p$, we see

$$\begin{aligned}|f(x)-f'(x)| &\le |f(x)-f(x')| + |f(x')-f'(x')| + |f'(x')-f'(x)| \\ &\le d_{X_p^n}(x,x') + \delta^{1-p} d_{X_1^n}(x,x') \\ &\le \delta + \delta^{1-p}\delta^p = 2\delta.\end{aligned}$$

Therefore,

$$\begin{aligned}\operatorname{diam}(f_*\mu_{X_p^n}; 1-\kappa) &\le \operatorname{diam}(f'_*\mu_{X_1^n}; 1-\kappa) + 4\delta \\ &\le n^{\frac{1}{2p}-\frac{1}{2}} \operatorname{ObsDiam}(X_1^n; -\kappa) + 4n^{\frac{1}{2p}}.\end{aligned}$$

By the arbitrariness of f, this completes the proof. □

Lemma 3.3. *Let X and Y be two mm-spaces. For any natural number n and real number p with $1 \le p \le +\infty$, we have*

$$\square(X_p^n, Y_p^n) \le n \cdot n! \, \square(X,Y).$$

Proof. We prove the lemma only in the case of $1 \le p < +\infty$. The proof for $p = +\infty$ is similar.

Note that $\square \le 1$ and that the lemma is trivial if $\square(X,Y) = 1$. Assume $\square(X,Y) < \varepsilon \le 1$. It suffices to prove that $\square(X_p^n, Y_p^n) \le n \cdot n! \, \varepsilon$. There are parameters $\varphi : I \to X$, $\psi : I \to Y$, and a subset $I_0 \subset X$ such that $\mathcal{L}^1(I_0) > 1-\varepsilon$ and

$$| \varphi^* d_X(s,t) - \psi^* d_Y(s,t) | < \varepsilon$$

for any $s,t \in I_0$. Define functions $\varphi_n : I^n \to X_p^n$ and $\psi_n : I^n \to Y_p^n$ by

$$\begin{aligned}\varphi_n(s_1,\dots,s_n) &:= (\varphi(s_1),\dots,\varphi(s_n)), \\ \psi_n(s_1,\dots,s_n) &:= (\psi(s_1),\dots,\psi(s_n))\end{aligned}$$

for $(s_1,\dots,s_n) \in I^n$. For any $s = (s_1,\dots,s_n), t = (t_1,\dots,t_n) \in I_0^n$, we have

$$\begin{aligned}&| \varphi_n^* d_{X_p^n}(s,t) - \psi_n^* d_{Y_p^n}(s,t) | \\ &= \left| \left(\sum_{i=1}^n d_X(\varphi(s_i),\varphi(t_i))^p \right)^{\frac{1}{p}} - \left(\sum_{i=1}^n d_Y(\psi(s_i),\psi(t_i))^p \right)^{\frac{1}{p}} \right| \\ &\le \left(\sum_{i=1}^n | \, d_X(\varphi(s_i),\varphi(t_i) - d_Y(\psi(s_i),\psi(t_i)) \, |^p \right)^{\frac{1}{p}} \\ &< n^{\frac{1}{p}} \varepsilon\end{aligned}$$

and

$$\mathcal{L}^n(I_0^n) \ge \mathcal{L}^1(I_0)^n > (1-\varepsilon)^n \ge 1 - n \cdot n! \, \varepsilon.$$

Since there is a Borel isomorphism from I to I^n that pushes $\mathcal{L}^1$ to $\mathcal{L}^n$ (see [4, (17.41)]), the proof is completed. □

Proof of Theorem 1.1. It suffices to prove the theorem only in the case where $\operatorname{diam}(X) = 1$, because of

$$\operatorname{ObsDiam}(X_p^n; -\kappa) = \operatorname{diam}(X) \operatorname{ObsDiam}((\operatorname{diam}(X)^{-1} X)_p^n; -\kappa)$$

for $\operatorname{diam}(X) > 0$. We assume $\operatorname{diam}(X) = 1$.

If $p = 1$, then [5, Corollary 1.7] implies

$$\alpha_{X_1^n}(r) \le \exp\left(-\frac{r^2}{8 \operatorname{diam}(X)^2 n}\right),$$

which together with Proposition 2.6(1) proves the theorem.

Assume $p > 1$. By [10, §4], there is a sequence of finite mm-spaces Y_k, $i = 1, 2, \ldots$, with diameter ≤ 1 that $\square$-converges to X. In addition, we may assume that the μ_{Y_k}-measure of each point in Y_k is an integral multiplication of $1/k$. In particular, Y_k consists of at most k points. Denoting by Z_k a k-regular mm-space, we see that $Y_k \prec Z_k$ for each k. By $(Y_k)_p^n \prec (Z_k)_p^n$, Lemma 3.2, and by the theorem for $p = 1$, we have

$$\begin{aligned}
\operatorname{ObsDiam}((Y_k)_p^n; -\kappa) &\le \operatorname{ObsDiam}((Z_k)_p^n; -\kappa) \\
&\le n^{\frac{1}{2p} - \frac{1}{2}} \operatorname{ObsDiam}((Z_k)_1^n; -\kappa) + 4 n^{\frac{1}{2p}} \\
&\le C_{\kappa,p}\, n^{\frac{1}{2p}}.
\end{aligned}$$

Lemma 3.3 implies that $(Y_k)_p^n$ $\square$-converges to X_p^n as $k \to \infty$. The limit formula (Theorem 2.11) proves

$$\operatorname{ObsDiam}(X_p^n; -\kappa) = \lim_{\varepsilon \to 0+} \liminf_{k \to \infty} \operatorname{ObsDiam}((Y_k)_p^n; -(\kappa+\varepsilon)) \le C_{\kappa,p}\, n^{\frac{1}{2p}}.$$

This completes the proof of the theorem. $\square$

Proof of Theorems 1.2 and 1.3. Take an arbitrary point $x_0 \in X$ and fix it. We define

$$f_n(x) := \sum_{i=1}^n d_X(x_0, x_i)^p$$

for $x = (x_1, x_2, \ldots, x_n) \in X^n$. Note that f_n is the sum of a sequence of independent and identically distributed random variables. The mean of f_n is

$$\int_{X^n} f_n \, d\mu_X^{\otimes n} = n E_p(x_0),$$

where $E_p(x_0) := \int_X d_X(x_0, \cdot)^p \, d\mu_X$. The variance of f_n is

$$\int_{X^n} (f_n - n E_p(x_0))^2 \, d\mu_X^{\otimes n} = n V_p(x_0),$$

where $V_p(x_0) := \int_X (d_X(x_0, \cdot)^p - E_p(x_0))^2 \, d\mu_X$. The central limit theorem tells us that

$$\text{(CLT)} \qquad \left(\frac{f_n - n E_p(x_0)}{\sqrt{n V_p(x_0)}}\right)_* \mu_X^{\otimes n} \to \gamma^1 \quad \text{weakly as } n \to \infty.$$

Take a real number κ with $0 < \kappa < 1/2$ and let

$$\begin{aligned}\alpha_n &:= \inf\{\, x \in \mathbb{R} \mid (f_n)_*\mu_X(\,-\infty, x\,] \geq \kappa\,\},\\ \beta_n &:= \sup\{\, x \in \mathbb{R} \mid (f_n)_*\mu_X[\,x, +\infty\,) \geq \kappa\,\},\\ c_n &:= \frac{\beta_n - \alpha_n}{\sqrt{nV_p(x_0)}}.\end{aligned}$$

Then, (CLT) proves that $\lim_{n\to\infty} c_n = 2\Phi^{-1}(\kappa)$. Letting

$$A_n := f_n^{-1}(\,-\infty, \alpha_n\,] \quad \text{and} \quad B_n := f_n^{-1}[\,\beta_n, +\infty\,),$$

we have

$$\mu_X^{\otimes n}(A_n), \mu_X^{\otimes n}(B_n) \geq \kappa.$$

Let us estimate $d_{X_p^n}(A_n, B_n)$ from below. For any points $x \in A_n$ and $y \in B_n$, we see that

$$\begin{aligned}c_n\sqrt{nV_p(x_0)} = \beta_n - \alpha_n &\leq |\, f_n(x) - f_n(y)\,|\\ &\leq \sum_{i=1}^n |\, d_X(x_0, x_i)^p - d_X(x_0, y_i)^p\,|\end{aligned}$$

and by $|a^p - b^p| \leq p \cdot \max\{a, b\}^{p-1}|a - b|$ for $a, b \geq 0$,

$$\begin{aligned}&\leq p\,\mathrm{diam}(X)^{p-1}\sum_{i=1}^n |\, d_X(x_0, x_i) - d_X(x_0, y_i)\,|\\ &\leq p\,\mathrm{diam}(X)^{p-1}\sum_{i=1}^n d_X(x_i, y_i)\\ &= p\,\delta\,\mathrm{diam}(X)^{p-1}\sum_{i=1}^n \frac{d_X(x_i, y_i)}{\delta}\end{aligned}$$

and by the assumption $\inf_{x\neq y} d_X(x, y) \geq \delta$,

$$\begin{aligned}&\leq p\,\delta\,\mathrm{diam}(X)^{p-1}\sum_{i=1}^n \frac{d_X(x_i, y_i)^p}{\delta^p}\\ &= p\,\delta^{1-p}\,\mathrm{diam}(X)^{p-1} d_{X_p^n}(x, y)^p,\end{aligned}$$

which implies

$$d_{X_p^n}(x, y) \geq \left(\frac{c_n\sqrt{V_p(x_0)}}{p}\right)^{\frac{1}{p}} \left(\frac{\delta}{\mathrm{diam}(X)}\right)^{1-\frac{1}{p}} n^{\frac{1}{2p}}.$$

In the case where $p = 1$, we do not need the assumption $\inf_{x\neq y} d_X(x, y) \geq \delta$ and obtain

$$d_{X_1^n}(x, y) \geq c_n\sqrt{nV_p(x_0)}.$$

Thus, if $0 < \kappa' < \kappa$, then

$$\begin{aligned}\mathrm{ObsDiam}(X_p^n; -\kappa') &\geq \mathrm{Sep}(X_p^n; \kappa, \kappa) \\ &\geq \left(\frac{c_n\sqrt{V_p(x_0)}}{p}\right)^{\frac{1}{p}} \left(\frac{\delta}{\mathrm{diam}(X)}\right)^{1-\frac{1}{p}} n^{\frac{1}{2p}}.\end{aligned}$$

This completes the proof. □

Proposition 3.4 ([2], [3, §3.$\frac{1}{2}$.42])**.** *Let X be a compact and connected Riemannian manifold with dimension ≥ 1, and let $0 < \kappa < 1$. We assume the measure μ_X to be the normalized volume measure on X, i.e., $\mu_X = \mathrm{vol} \,/\, \mathrm{vol}(X)$, where* vol *is the volume measure on X. Then, the sequence $\{\mathrm{ObsDiam}(X_2^n; -\kappa)\}_{n=1}^{\infty}$ is bounded from above and bounded away from zero.*

Proof. Since a natural projection $X_2^n \to X$ induces $X \prec X_2^n$, we have the lower estimate

$$0 < \mathrm{ObsDiam}(X; -\kappa) \leq \mathrm{ObsDiam}(X_2^n; -\kappa).$$

Denote by $\lambda_1(X)$ the first non-zero eigenvalue of the Laplacian on X. By $\lambda_1(X_2^n) = \lambda_1(X)$, we have the upper estimate

$$\mathrm{ObsDiam}(X_2^n; -\kappa) \leq \frac{2\sqrt{2}}{\sqrt{\lambda_1(X)\kappa}}$$

(see [10, Corollary 2.39]). This proves the proposition. □

Combining Proposition 3.4 with Theorem [9, Theorem 1.2] implies

Corollary 3.5. *If X is a compact and connected Riemannian manifold with dimension ≥ 1, then $\{X_2^n\}_{n=1}^{\infty}$ has the phase transition property with critical scale order ~ 1.*

References

[1] D. Amir and V. D. Milman, *Unconditional and symmetric sets in n-dimensional normed spaces*, Israel J. Math. **37** (1980), no. 1-2, 3–20.

[2] M. Gromov and V. D. Milman, *A topological application of the isoperimetric inequality*, Amer. J. Math. **105** (1983), no. 4, 843–854.

[3] M. Gromov, *Metric structures for Riemannian and non-Riemannian spaces*, Reprint of the 2001 English edition, Modern Birkhäuser Classics, Birkhäuser Boston Inc., Boston, MA, 2007. Based on the 1981 French original; With appendices by M. Katz, P. Pansu and S. Semmes; Translated from the French by Sean Michael Bates.

[4] A. S. Kechris, *Classical descriptive set theory*, Graduate Texts in Mathematics, vol. 156, Springer-Verlag, New York, 1995.

[5] M. Ledoux, *The concentration of measure phenomenon*, Mathematical Surveys and Monographs, vol. 89, American Mathematical Society, Providence, RI, 2001.

[6] P. Lévy, *Problèmes concrets d'analyse fonctionnelle. Avec un complément sur les fonctionnelles analytiques par F. Pellegrino*, Gauthier-Villars, Paris, 1951 (French). 2d ed.

[7] V. D. Milman, *The heritage of P. Lévy in geometrical functional analysis*, Astérisque **157-158** (1988), 273–301. Colloque Paul Lévy sur les Processus Stochastiques (Palaiseau, 1987).
[8] V. D. Milman and G. Schechtman, *Asymptotic theory of finite-dimensional normed spaces*, Lecture Notes in Mathematics, vol. 1200, Springer-Verlag, Berlin, 1986. With an appendix by M. Gromov.
[9] R. Ozawa and T. Shioya, *Limit formulas for metric measure invariants and phase transition property*. preprint.
[10] T. Shioya, *Metric measure geometry–Gromov's theory of convergence and concentration of metrics and measures*. preprint.
[11] M. Talagrand, *Concentration of measure and isoperimetric inequalities in product spaces*, Inst. Hautes Études Sci. Publ. Math. **81** (1995), 73–205.

Mathematical Institute, Tohoku University, Sendai 980-8578, JAPAN